



Tests and Turnaround Times

Test	Turnaround Time*
Cardiovascular disease/dyslipidemia	
Apo E Genotype Test (for type III hyperlipoproteinemia)	1 day
Celiac disease	
Celiac Disease DNA Test	1 day
Celiac Disease Antibody Panel tTG-IgA, EMA-IgA, Gliadin IgA and IgG, Total IgA	2-3 days
Endomysial (EMA) IgA Antibody	2-3 days
Gliadin IgA Antibody	2-3 days
Gliadin IgG Antibody	2-3 days
Gliadin IgA and IgG Antibodies	2-3 days
Reticulin IgA Antibody	2-3 days
Reticulin Total Antibodies	2-3 days
Tissue Transglutaminase (tTG) IgA Antibody	2-3 days
Tissue Transglutaminase (tTG) IgG Antibody	2-3 days
Tissue Transglutaminase (tTG) IgA and IgG Antibodies	2-3 days
Total IgA	2-3 days
Cystic fibrosis	
Cystic Fibrosis DNA Test – Carrier screening and diagnostic testing	1 day
Cystic Fibrosis DNA Test – Prenatal testing (CVS or amnio, direct or cultured)	3 days
Fragile X syndrome and related testing	
Fragile X DNA Test – Carrier and diagnostic testing	7-11 days
Fragile X DNA Test – Prenatal testing (CVS or amnio, direct or cultured)	7-11 days
Fragile X Protein (FMRP) Test	2-3 weeks
Fragile X DNA/Standard Chromosome Analysis Combined Test	7-14 days
FRAXE DNA Test	10-14 days
FXTAS DNA Test (Fragile X-Associated Tremor/Ataxia Syndrome)	7-11 days
Hemochromatosis	
Hemochromatosis DNA Test	1 day
HLA-B27 (ankylosing spondylitis)	
HLA-B27 DNA Test	1 day

* Turnaround time represents business days.



Tests and Turnaround Times

Test

Turnaround Time*

Hypercoagulability

Factor V Leiden DNA Test	1 day
Prothrombin (Factor II) DNA Test	1 day
Factor V Leiden/Prothrombin (Factor II) Combined DNA Test	1 day
Factor V R2 DNA Test	1 day
MTHFR DNA Test	1 day
Inherited Hypercoagulability Panel A (for patients not on coumadin therapy) [†]	2-3 days
Factor V Leiden DNA Test, Prothrombin (Factor II) DNA Test, Antithrombin III Activity, Protein C Activity, Protein S Activity	
Inherited Hypercoagulability Panel B (for patients on coumadin therapy) [†]	4-6 days
Factor V Leiden DNA Test, Prothrombin (Factor II) DNA Test, Antithrombin III Activity, Protein C/Factor IX Antigen Ratio, Protein S/Factor IX Antigen Ratio	
High Risk Pregnancy Hypercoagulability Panel [†]	7 days
Factor V Leiden DNA Test, Prothrombin (Factor II) DNA Test, Antithrombin III Activity, Protein C Activity, Protein S Activity, Antiphospholipid Antibody Syndrome Tests (Lupus Anticoagulant and Anticardiolipin Antibodies IgG, IgM, and IgA),	
Anti-β ₂ Glycoprotein I Antibody Test (IgG, IgM, and IgA)	3-5 days
Homocysteine (serum)	5-6 days

[†] Tests in panels can be ordered individually. Call for details.

Narcolepsy

Narcolepsy DNA Test	2 days
---------------------	--------

Periodontal disease

PST® Genetic Test for Susceptibility to Periodontal Disease	1-2 days
---	----------

PST® is a registered service mark of Interleukin Genetics, Inc.
PST® genotyping is performed under a license from Interleukin Genetics, Inc.

Tay-Sachs disease

Tay-Sachs Disease DNA Test – Carrier screening and diagnostic testing	1 day
Tay-Sachs Disease DNA Test – Prenatal testing (1278insTATC, 1421+1G→C, and G805A) (CVS or amnio, direct or cultured)	3 days
Tay-Sachs Enzyme Test	3-5 days

* Turnaround time represents business days.

